

WHAT IS CLAIMED IS:

1. An isolated or recombinant polynucleotide encoding an antigenic polypeptide comprising at least 17
5 contiguous amino acids from the mature polypeptide from SEQ ID NO: 2 or 4.
2. The polynucleotide of Claim 1, encoding a mature polypeptide from SEQ ID NO: 2 or 4.
- 10 3. The polynucleotide of Claim 1, which hybridizes at 55° C, less than 500 mM salt, and 50% formamide to the coding portions of SEQ ID NO: 1 or 3.
- 15 4. The polynucleotide of Claim 3, comprising at least 35 contiguous nucleotides of the coding portion of SEQ ID NO: 1 or 3.
5. An expression vector comprising the
20 polynucleotide of Claim 1.
6. A host cell containing the expression vector of Claim 5, including a eukaryotic cell.
- 25 7. A method of making an antigenic polypeptide comprising expressing a recombinant polynucleotide of Claim 1.
8. A method for forming a duplex with a
30 polynucleotide of Claim 1, comprising contacting said polynucleotide with a probe that hybridizes, under stringent conditions, to at least 25 contiguous nucleotides of the coding portion of SEQ ID NO: 1 or 3; thereby forming said duplex.

9. A kit for the detection of a polynucleotide of Claim 1, comprising a polynucleotide that hybridizes, under stringent hybridization conditions, to at least 17 contiguous nucleotides of a polynucleotide of Claim 1.

5 10. The kit of claim 9, wherein said probe is detectably labeled.

11. A binding compound comprising an antibody binding site which specifically binds to at least 17 contiguous
10 amino acids from SEQ ID NO: 2 or 4.

12. The binding compound of Claim 11, wherein:

a) said antibody binding site is:

- 15 1) specifically immunoreactive with a polypeptide of SEQ ID NO: 2 or 4;
2) raised against a purified or recombinantly produced human IL-B50 protein; or
3) in a monoclonal antibody, Fab, or F(ab)2; or

b) said binding compound is:

- 20 1) an antibody molecule;
2) a polyclonal antiserum;
3) detectably labeled;
4) sterile; or
5) in a buffered composition.

25 13. A method using the binding compound of Claim 11, comprising contacting said binding compound with a biological sample comprising an antigen, wherein said contacting results in formation of a binding
30 compound:antigen complex.

14. The method of Claim 13, wherein said biological sample is from a human, and wherein said binding compound is an antibody.

35

15. A detection kit comprising said binding compound of Claim 12, and:

- a) instructional material for the use of said binding compound for said detection; or
- 5 b) a compartment providing segregation of said binding compound.

16. A substantially pure or isolated antigenic polypeptide, which binds to said binding composition of Claim 11, and further comprises at least 17 contiguous amino
10 acids from SEQ ID NO: 2 or 4.

17. The polypeptide of Claim 16, which:

- a) comprises at least a fragment of at least 25 contiguous amino acid residues from a primate
15 IL-B50 protein;
- b) is a soluble polypeptide;
- c) is detectably labeled;
- d) is in a sterile composition;
- e) is in a buffered composition;
- 20 f) binds to a cell surface receptor;
- g) is recombinantly produced; or
- h) has a naturally occurring polypeptide sequence.

18. The polypeptide of Claim 17, which comprises at
25 least 17 contiguous amino acids of SEQ ID NO: 2 or 4.

19. A method of modulating physiology or development of a cell or tissue culture cells comprising contacting said cell with an agonist or antagonist of a primate IL-B50.
30

20. The method of Claim 19, wherein:

- a) said contacting is in combination with an agonist or antagonist of IL-7; or
- 35 b) said contacting is with an antagonist, including binding composition comprising an antibody binding site which specifically binds an IL-B50.